

## **BIOPSIA HEPÁTICA EN HEPATOPATÍAS DIFUSAS: indicaciones, técnica, limitaciones y resultados. Alternativas endovasculares.**

### **Bibliografía:**

- Terry R. Risks of needle biopsy of liver. Br.M.J. 1952.1:1102-1105.
- Zamcheck N. Liver biopsy.II. The risk of needle biopsy. New E.J.M. ,1953 vol.249 N°26 :1062-1069.
- Piccinino F et al. Complications following percutaneous liver biopsy. A multicenter retrospective study on 68,276 biopsies. J.Hepatol 1986;2:165-173.
- Chuah S.Y. et al. A nationwide survey of liver biopsy.- Is there a need to increase resources, manpower and training?. Hepato-Gastroenterol.41 (1994) 4-8.
- Sugano S. et al . Incidence of ultrasound-detected intrahepatic hematomas due to Tru-cut needle liver biopsy. Dig.Dis.Sci. Vol 36.N°9 (1991) 1229-1233.
- Hederstrom E. et al. Liver biopsy complications monitored by ultrasound. J.Hepatol.(1989) 8: 94-98.
- Minuk J.I. et al Prospective study of the incidence of ultrasound detected intrahepatic and subcapsular hematomas in patients randomized to 6 or 24 hours of bed rest after percutaneous liver biopsy. Gastroenterology 92:290-293, 1987.
- Caturelli E. Biopsia epatica percutanea . en Quaderni di ecografia operativa. Poletto Edizioni. Milano feb.1997.
- Buscarini E- Di Stasi M. Complications of abdominal interventional ultrasound. Poletto edizioni. Milano 1996.
- Joly JP, Khouani S, Decrombeque C et al : la ponction-biopsie hépatique écouguidée pourrait remplacer la ponction-biopsie à l'aveugle dans les hépatopathies diffuses. Gstroenterol.Clin.Biol. 19: 703-706, 1995.
- Lindor KD, Bru C, Jorgensen RA et al: The role of ultrasonography and automatic-needle biopsy in outpatient percutaneous liver biopsy. Hepatology, 23: 1079-1083, 1996.
- Caturelli E, Giacobbe A, Facciorusso D et al: Percutaneous biopsy in diffuse liver disease: increasing diagnostic yield and decreasing complication rate by routine ultrasound assessment of puncture site. Am.J.Gastroenterol, 1996,91: 1318-1321.
- Pasha T, Gabriel S, Therneau T , Dikson ER, Lindor KD: Cost-effectiveness of US-guided liver biopsy. Hepatology 1998, 27(5)1220-6.
- Younossi ZM, Teran JC, Ganiats TG, Carey WD : US-guided liver biopsy for parenchymal liver disease: an economic analysis. Dig Dis Sci Jan 1998, 43(1)46-50.
- Farrell RJ, Smiddy PF, Pilkington RM, Tobin AA,Mooney EE et al : Guided versus blind liver biopsy for chronic hepatitis C :clinical benefits and costs . J Hepatol 1999;30:580-587.
- Candranel JF,Rufat P, Degos F : Practices of liver biopsy in France: results of a prospective nationwide survey. Hepatology Vol 32 n°3 2000: 477-81.
- McGill DB,: Liver biopsy : when, how,by whom and where? Current Gastroenterology 2000; 1:19-23.
- Jackson JE, Adam, A, Allison DJ. : Transjugular and plugged liver biopsies. Baillieres Clin Gatrienterol 1992,6(2):245-58.
- Mendoza J,Gomez\_Dominguez E, Moreno -Otero R. Elastografía de transición (fibrosan), un nuevo método no invasivo en la valoración de la fibrosis hepática. Med Clin (Barc) 2006,Febrero 18;126(6):220-2.