

## **Tècniques d'imatge a urologia pediàtrica .**

### **Bibliografia:**

1. T Berrocal, I Pinilla, J Gutierrez , C Prieto, Lde Pablo, ML del Hoyo. Mild hydronephrosis in newborns and infants: can ultrasound predict the presence of vesicoureteral reflux. *Pediatric Nephrol* 2007; 22:91-96.
2. M Riccabona and R Fötter. Reorientation and future trends in pediatric uroradiology. *Pediatric Radiol* 2004;34:295-301.
3. T Berrocal, F Gaya, A Arjonilla, G Lonergan. Vesicoureteral reflex:diagnosis and grading with echo-enhanced cystosonography versus voiding cystourethrography. *Radiology* 2001;221:359-365.
4. T Berrocal, F Gaya, A Arjonilla. Vesicoureteral Reflux : Can the urethra be adequately assessed by using contrast-enhanced voiding US of the bladder?. *Radiology* 2005; 234:235-241.
5. K Darge, RT Moeller, A Trusen, F Butter,N Gordjani, H Riedmiller. Diagnosis of vesicoureteric reflux with low-dose contrast-enhanced harmonic ultrasound imaging. *Pediatric Radiol* 2005; 35: 73-78.
6. RL Lebowitz, H Olbing, KV Parkkulainen, JM Smellie and TE Tamminen-Möbius. International system of radiographic grading of vesicoureteric reflux. *Pediatric Radiol* (1985) 15:105-109