

## Bibliografía

1. ACCF/ACR/SCCT/SCMR/ASNC/NASCI/SCAI/SIR 2006 appropriateness criteria for cardiac computed tomography and cardiac magnetic resonance imaging: a report of the American College of Cardiology Foundation Quality Strategic Directions Committee Appropriateness Criteria Working Group, American College of Radiology, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, American Society of Nuclear Cardiology, North American Society for Cardiac Imaging, Society for Cardiovascular Angiography and Interventions, and Society of Interventional Radiology. Hendel, R.C.; Patel, M.R.; Kramer, C.M.; Poon, M.; Hendel, R.C.; Carr, J.C.; Gerstad, N.A.; Gillam, L.D.; Hodgson, J.M.; Kim, R.J.; Kramer, C.M.; Lesser, J.R.; Martin, E.T.; Messer, J.V.; Redberg, R.F.; Rubin, G.D.; Rumsfeld, J.S.; Taylor, A.J.; Weigold, W.G.; Woodard, P.K.; Brindis, R.G.; Hendel, R.C.; Douglas, P.S.; Peterson, E.D.; Wolk, M.J.; Allen, J.M.; Patel, M.R. *J. Am Coll. Cardiol.* 2006;48:1475-1497.
2. Comprehensive cardiac CT study: evaluation of coronary arteries, left ventricular function, and myocardial perfusion--is it possible? Cury, R.C.; Nieman, K.; Shapiro, M.D.; Nasir, K.; Cury, R.C.; Brady, T.J. *J. Nucl. Cardiol.* 2007;14:229-243.
3. The salvaged area at risk in reperfused acute myocardial infarction as visualized by cardiovascular magnetic resonance. Friedrich, M.G.; Abdel-Aty, H.; Taylor, A.; Schulz-Menger, J.; Messroghli, D.; Dietz, R. *J. Am Coll. Cardiol.* 2008;51:1581-1587.
4. MRI delayed contrast enhancement to irreversible injury, infarct age, and contractile function. Kim RJ, Fieno DS, Parrish TB, Harris K, Chen EL, Simonetti O, Bundy J, Finn JP, Klocke FJ, Judd R. *Circulation*, 1999;19:1992-2002.
5. Assessment of myocardial viability with contrast-enhanced magnetic resonance imaging: comparison with positron emission tomography. Klein, C.; Nekolla, S.G.; Bengel, F.M.; Momose, M.; Sammer, A.; Haas, F.; Schnackenburg, B.; Delius, W.; Mudra, H.; Wolfram, D.; Schwaiger, M. *Circulation*, 2002;105:162-167.
6. The use of contrast-enhanced magnetic resonance imaging to identify reversible myocardial dysfunction. Kim, R.J.; Wu, E.; Rafael, A.; Chen, E.L.; Parker, M.A.; Simonetti, O.; Klocke, F.J.; Bonow, R.O.; Judd, R.M. *N Engl. J. Med.* 2000;343:1445-1453.
7. Contrast-enhanced MRI and routine single photon emission computed tomography (SPECT) perfusion imaging for detection of subendocardial myocardial infarcts: an imaging study. Wagner, A.; Mahrholdt, H.; Holly, T.A.; Elliott, M.D.; Regenfus, M.; Parker, M.; Klocke, F.J.; Bonow, R.O.; Kim, R.J.; Judd, R.M. *Lancet*, 2003; 361:374-379.