

LESIONES FOCALES HEPÁTICAS: UN ENFOQUE CLÍNICO Y RADIOLÓGICO

Bibliografía:

1. Baron RL. Understanding and optimizing use of contrast material for CT of the liver. *AJR* 1994;163:323-331
2. Freeny PC, Baron RL, Teefey SA. Hepatocellular carcinoma: reduced frequency of typical findings with dynamic contrast-enhanced CT in non-asian population. *Radiology* 1992;182:143-148
3. Stevens WR, Johnson CD, Stephens DH, Batts KP. CT findings in hepatocellular carcinoma: correlation of tumor characteristics with causative factors, tumor size and histologic grade. *Radiology* 1994;191:513
4. Hanafusa K, Ohashi I, Himeno Y, Suzuki S, Shibuya H. Hepatic hemangiomas: findings with two phase CT. *Radiology* 1995;196:465-469
5. Honda H, Onitsuka H, Yashumori K et al. Intrahepatic peripheral colangiocarcinoma: two-phased dynamic incremental CT and pathologic correlation. *J Comput Assist Tomogr* 1993;17:3907-402
6. Kim TK, Choi BI, Han JK, Jang HJ, Cho SG, Han MC. Peripheral cholangiocarcinoma of the liver: two-phase spiral CT findings. *Radiology* 1997;204:539-543
7. Freeny PC, Marks WM. Hepatic hemangiomas: dynamic bolus CT. *AJR* 1986;147:711-719
8. Leslie DF, Johnson CD, Johnson CM, Ilstrup DM, Harmsen WS. Distinction between cavernous hemangiomas of the liver and hepatic metastases on CT: value of contrast enhancement patterns. *AJR* 1995;164:625-629
9. Van Leuveen MS, Noordzij J, Feldberg MAM, Hennipman A, Doorneward H. Focal liver lesions: characterization with triphasic spiral CT. *Radiology* 1996;201:327-336
10. Jones EC, Chezmar JL, Nelson RL, Bernardino ME. The frequency and significance of small (>15 mm) hepatic lesions detected by CT. *AJR* 1992;158:535-539
11. Schwartz L *Radiology* 1999;210:71-74
12. Buetow PC, Pantongrag-Brown L, Buck JL, Ros PR, Goodman ZD. Focal nodular hyperplasia: pathologic radiologic correlation. *Radiographics* 1996;16:369-388