

TRACTAMENT PERCUTANI DE LA PATOLOGIA BILIAR: PATOLOGIA BENIGNA I MALIGNA

Bibliografia:

1. B. Topal, et al. The value of magnetic resonance cholangiopancreatography in predicting common bile duct stones in patients with gallstone disease. *British Journal of Surgery* 2003; 90: 42-47
2. J. Lopera, et.al. Biliary Obstruction: Use of MR Cholangiography to Define the Extent of Biliary Ductal Involvement and Plan Percutaneous Interventions. *220:90 Radiology* 2001
3. Van Delden OM, Laméris JS. Percutaneous drainage and stenting for palliation of malignant bile duct obstruction. *Eur Radiol* 2008 Mar;18(3):448-56.
4. V Piñol, A Castells, J. Bordas, I. Real, J. Llach, X. Montanyà, F. Feu, S. Navarro. Percutaneous Self-expanding Metal Stents versus Endoscopic Polyethylene Endoprosthesis for Treating Malignant Biliary Obstruction: Randomized Clinical Trial. *Radiology* 2002; 225:27-34
6. N. Fidelman, et al. Hepatic Arterial Injuries after Percutaneous Biliary Interventions in the Era of Laparoscopic Surgery and Liver Transplantation: Experience with 930 Patients. *Radiology* 247: Number 3 June 2008
7. Schumacher B. et al.. Long-term follow-up of percutaneous transhepatic therapy (PTT) in patients with definite benign anastomotic strictures after hepaticojejunostomy. *Endoscopy*. 2001
8. Draganov P. Long-term outcome in patients with benign biliary strictures treated endoscopically with multiple stents. *Gastrointest Endosc* 2002 May;55(6):680-6